

ON CERTAIN CLASSES OF HARMONIC FUNCTIONS DEFINED BY THE FRACTIONAL DERIVATIVES

M. ESHAGHI GORDJI, S. SHAMS AND A. EBADIAN

ABSTRACT. In this paper we have introduced two new classes $\mathcal{HM}(\beta, \lambda, k, \nu)$ and $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ of complex valued harmonic multivalent functions of the form $f = h + \bar{g}$, satisfying the condition

$$\operatorname{Re} \left\{ (1 - \lambda) \frac{\Omega^\nu f}{z} + \lambda(1 - k) \frac{(\Omega^\nu f)'}{z'} + \lambda k \frac{(\Omega^\nu f)''}{z''} \right\} > \beta, \quad (z \in \mathcal{D})$$

where h and g are analytic in the unit disk $\mathcal{D} = \{z : |z| < 1\}$. A sufficient coefficient condition for this function in the class $\mathcal{HM}(\beta, \lambda, k, \nu)$ and a necessary and sufficient coefficient condition for the function f in the class $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ are determined. We investigate inclusion relations, distortion theorem, extreme points, convex combination and other interesting properties for these families of harmonic functions.

1. INTRODUCTION

Let u, v be real harmonic function in a simply connected domain Ω , then the continuous function $f = u + iv$ defined in Ω is said to be harmonic in Ω . If $f = u + iv$ is harmonic in Ω then there exist analytic functions G, H such that $u = \operatorname{Re} G$ and $v = \operatorname{Im} H$, therefor $f = u + iv = h + \bar{g}$ where $h = \frac{G+H}{2}$, $\bar{g} = \frac{\overline{G-H}}{2}$ and we call h and g analytic part and co-analytic part of f respectively. The jacobian of f is given by $J_f|z| = |h'(z)|^2 - |g'(z)|^2$, also we show by $w(z)$ the dilatation function for f and define $w(z) = \frac{g'(z)}{h'(z)}$. Lewy [6], Clunie and Small [3] have showed that the mapping $z \rightarrow f(z)$ is sense preserving and injective in Ω if and only if $J_f|z| > 0$ in Ω . The function $f = h + \bar{g}$ is said to be univalent in Ω if the mapping $z \rightarrow f(z)$ is sense preserving and injective in Ω . Denote by $\mathcal{H}$ the class of all harmonic functions $f = h + \bar{g}$ that are univalent and sense preserving in the open unit disk $\mathcal{D}$ where

$$(1.1) \quad h(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad g(z) = \sum_{n=1}^{\infty} b_n z^n \quad |b_1| < 1.$$

With normalization conditions $f(0) = 0$, $f_z(0) = 1$ where $f_z(0)$ denotes the partial derivative of $f(z)$ at $z = 0$. In case $g = 0$ this class reduces to the class of $\mathcal{S}$ consisting of all analytic univalent functions.

Definition 1.1. (See [7] and [9]) Let the function $f(z)$ be analytic in a simply-connected region of the z -plane containing the origin. The fractional derivative of f of order ν is defined by

$$D_z^\nu f(z) = \frac{1}{\Gamma(1-\nu)} \frac{d}{dz} \int_0^1 \frac{f(\zeta)}{(z-\zeta)^\nu} d\zeta, \quad 0 \leq \nu < 1$$

where the multiplicity of $(z-\zeta)^\nu$ is removed by requiring $\log(z-\zeta)$ to be real when $z-\zeta > 0$.

2000 *Mathematics Subject Classification.* Primary 30C45; Secondary 30C80.

Key words and phrases. Multivalent, Harmonic, Convex, Starlike, Convolution and Fractional derivatives.

Making use of fractional derivative and its known extensions involving fractional derivatives and fractional integrals, Owa and Srivastava [8] introduced the operator $\Omega_z^\nu : \mathcal{A}_0 \longrightarrow \mathcal{A}_0$ defined by

$$\Omega_z^\nu f(z) := \Gamma(2 - \nu) z^\nu D_z^\nu f(z) \quad \nu \neq 2, 3, 4, \dots$$

where $\mathcal{A}_0$ denote the class of functions which are analytic in the unit disk $\mathcal{D}$, satisfying normalization conditions $f(0) = f'(0) - 1 = 0$.

It is easy to see that

$$\Omega_z^\nu f(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(2 - \nu)\Gamma(n + 1)}{\Gamma(n + 1 - \nu)} a_n z^n, \quad f \in \mathcal{A}_0$$

Definition 1.2. Suppose that $f = h + \overline{g}$ where h and g are in (1.1), define $\Omega_z^\nu f(z) = \Omega_z^\nu h(z) + \overline{\Omega_z^\nu g(z)}$.

Then we obtain

$$\Omega_z^\nu f(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(2 - \nu)\Gamma(n + 1)}{\Gamma(n + 1 - \nu)} a_n z^n + \sum_{n=1}^{\infty} \frac{\Gamma(2 - \nu)\Gamma(n + 1)}{\Gamma(n + 1 - \nu)} b_n \overline{z}^n.$$

By making use of Definition 1.2, we introduce a new class of harmonic univalent functions in the unit disk $\mathcal{D}$ as in definition 1.3.

Definition 1.3. Let $\mathcal{HM}(\beta, \lambda, k, \nu)$ ($0 \leq k \leq 1$, $0 < \beta \leq 1$, $0 \leq \lambda$, $0 \leq \nu < 1$) be the class of functions $f \in \mathcal{H}$ satisfying the following inequality:

$$\operatorname{Re} \left\{ (1 - \lambda) \frac{\Omega^\nu f}{z} + \lambda(1 - k) \frac{(\Omega^\nu f)'}{z'} + \lambda k \frac{(\Omega^\nu f)''}{z''} \right\} > \beta, \quad (z = re^{i\theta})$$

where

$$z' = \frac{\partial}{\partial \theta} (re^{i\theta}), \quad z'' = \frac{\partial}{\partial \theta} (z'),$$

and

$$(\Omega^\nu f(z))' = \frac{\partial}{\partial \theta} (\Omega^\nu f(z)) = iz(\Omega^\nu h(z))' - i\overline{z(\Omega^\nu g(z))'},$$

$$(\Omega^\nu f(z))'' = \frac{\partial}{\partial \theta} (\Omega^\nu f(z))' = -z(\Omega^\nu h(z))' - z^2(\Omega^\nu h(z))'' - \overline{z(\Omega^\nu g(z))'} - \overline{z^2(\Omega^\nu g(z))''},$$

also we denote by $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ the subclass of $\mathcal{HM}(\beta, \lambda, k, \nu)$ consisting of functions $f = h + \overline{g}$ such that

$$(1.2) \quad h(z) = z - \sum_{n=2}^{\infty} |a_n| z^n, \quad g(z) = \sum_{n=1}^{\infty} |b_n| z^n, \quad |b_1| < 1.$$

In [9] H. M. Srivastava and S. Owa investigated this class with $D^\nu f(z)$ instead of $\Omega^\nu f(z)$ where $D^\nu f(z)$ is the Ruscheweyh derivative of f , for p -valent harmonic functions. This class in special cases involve the works studied by the previous authors such as Bhoosnurmath and Swamay [2], Ahuja and Jahangiri [1,5].

In this paper the coefficient inequalities for the classes $\mathcal{HM}(\beta, \lambda, k, \nu)$ and $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ are obtained also some other interesting properties of these classes are investigated.

2. COEFFICIENT BOUNDS

In the first theorem we give the sufficient condition for $f \in \mathcal{H}$ to be in the class $\mathcal{HM}(\beta, \lambda, k, \nu)$.

Theorem 2.1. Let $f \in \mathcal{H}$, and

$$\sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| < 1 - \beta,$$

where

$$(2.1) \quad \phi(n, k, \lambda, \nu) := \frac{[1 + \lambda(n-1)(1+nk)]\Gamma(n+1)\Gamma(2-\nu)}{\Gamma(n+1-\nu)},$$

and

$$(2.2) \quad \psi(n, k, \lambda, \nu) := \frac{[1 - \lambda(n+1)(1-nk)]\Gamma(n+1)\Gamma(2-\nu)}{\Gamma(n+1-\nu)},$$

then $f \in \mathcal{HM}(\beta, \lambda, k, \nu)$. The result is sharp for the function $f(z)$ given by

$$\begin{aligned} f(z) &= z + \sum_{n=2}^{\infty} \frac{\gamma^n \Gamma(n+1-\nu) z^n}{[1 + \lambda(n-1)(1+nk)]\Gamma(n+1)\Gamma(2-\nu)} \\ &\quad + \sum_{n=1}^{\infty} \frac{\delta^n \Gamma(n+1-\nu)}{[1 - \lambda(n+1)(1-nk)]\Gamma(n+1)\Gamma(2-\nu)} \bar{z}^n \end{aligned}$$

where $\sum_{n=2}^{\infty} |\gamma_n| + \sum_{n=1}^{\infty} |\delta_n| = 1 - \beta$.

Proof. Suppose

$$E(z) = (1-\lambda) \frac{\Omega^\nu f(z)}{z} + \lambda(1-k) \frac{(\Omega f(z))'}{z'} + \lambda k \frac{(\Omega f(z))''}{z''}.$$

It suffices to show that $|1 - \beta + E(z)| \geq |1 + \beta - E(z)|$. A simple calculation by substituting for h and g in $E(z)$ shows

$$\begin{aligned} E(z) &= 1 + \sum_{n=2}^{\infty} \frac{[1 + \lambda(n-1)(1+nk)]\Gamma(n+1)\Gamma(2-\nu)}{\Gamma(n+1-\nu)} a_n z^{n-1} \\ &\quad + \sum_{n=1}^{\infty} \frac{[1 - \lambda(n+1)(1-nk)]\Gamma(n+1)\Gamma(2-\nu)}{\Gamma(n+1-\nu)} b_n \frac{\bar{z}^n}{z}, \end{aligned}$$

Considering (2.1) and (2.2) we have

$$\phi(n, k, \lambda, \nu) = n(n-1)[1 + \lambda(n-1)(1+nk)]B(n-1, 2-\nu),$$

and

$$\psi(n, k, \lambda, \nu) = n(n-1)[1 - \lambda(n+1)(1-nk)]B(n-1, 2-\nu),$$

where $B(\alpha, \beta) = \int_0^1 t^{\alpha-1}(1-t)^{\beta-1}dt = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is the familiar Beta function. Then we obtain

$$E(z) = 1 + \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) a_n z^{n-1} + \sum_{n=1}^{\infty} \psi(k, n, \lambda, \nu) b_n \frac{\bar{z}^n}{z}.$$

Now we have

$$\begin{aligned}
& |1 - \beta + E(z)| - |1 + \beta - E(z)| \\
&= |2 - \beta + \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) a_n z^{n-1} + \sum_{n=1}^{\infty} \psi(n, k, \lambda, \nu) b_n \frac{\bar{z}^n}{z}| \\
&\quad - |\beta - \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) a_n z^{n-1} - \sum_{n=1}^{\infty} \psi(n, k, \lambda, \nu) b_n \frac{\bar{z}^n}{z}| \\
&\geq 2 - \beta + \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| |z|^{n-1} + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| \frac{\bar{z}^n}{z} \\
&\quad - \beta - \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| |z|^{n-1} - \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| \frac{\bar{z}^n}{z} \\
&= 2 - 2\beta - 2 \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| |z|^{n-1} - 2 \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| \frac{\bar{z}^n}{z} \\
&> 2 - 2\beta - 2 \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| |z|^{n-1} - 2 \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| \frac{\bar{z}^n}{z} \\
&\geq 0,
\end{aligned}$$

and the proof is complete. $\square$

In our next theorem we obtain the necessary and sufficient coefficients condition for the $f \in \mathcal{H}$ to be in $\overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu)$.

Theorem 2.2. *Let $f \in \mathcal{H}$ then $f \in \overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu)$ if and only if*

$$(2.3) \quad \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| < 1 - \beta.$$

Proof. Since $\overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu) \subset \mathcal{H}\mathcal{M}(\beta, \lambda, k, \nu)$ then the "if" part of theorem follows from Theorem 2.1, for "only if" part we show that if the condition (2.3) dose not hold then $f \notin \overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu)$. Let $f \in \overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu)$ then we have

$$\begin{aligned}
0 &\leq \operatorname{Re} \left\{ (1 - \lambda) \frac{\Omega^\nu f(z)}{z} + \lambda(1 - k) \frac{(\Omega^\nu f(z))'}{z'} + \lambda k \frac{(\Omega^\nu f(z))''}{z''} - \beta \right\} \\
&= \operatorname{Re} \left\{ 1 - \beta - \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) a_n z^{n-1} - \sum_{n=1}^{\infty} \psi(n, k, \lambda, \nu) b_n \frac{\bar{z}^n}{z} \right\}.
\end{aligned}$$

This inequality holds for all values of z for which $|z| = r < 1$ so we can choose the values of z on positive real axis such that $0 \leq z = r < 1$ therefore we get the followin inequality

$$0 \leq 1 - \beta - \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| r^{n-1} - \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| r^{n-1}.$$

Now by letting $r \rightarrow 1^-$ we have

$$(2.4) \quad 0 \leq 1 - \beta - \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| - \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n|.$$

If the condition (2.3) dose not hold then the right hand of (2.4) is negative for r sufficiently close to 1. Thus there exists a $z_0 = r_0 \in (0, 1)$ for which the right hand of (2.4) is negative. This contradicts the required condition for $f \in \overline{\mathcal{H}\mathcal{M}}(\beta, \lambda, k, \nu)$ and so the proof is complete. $\square$

Putting $\lambda = 0$ in Theorem 2.2 we get:

Corollary 2.1. $f \in \overline{\mathcal{HM}}(\beta, 0, k, \nu) = \left\{ f : \operatorname{Re} \left(\frac{\Omega^\nu f(z)}{z} \right) > \beta \right\}$ if and only if

$$\sum_{n=1}^{\infty} n(n-1)B(n-1, 2-\nu)|a_n| + \sum_{n=1}^{\infty} n(n-1)B(n-1, 2-\nu)|b_n| < 1-\beta.$$

Putting $\lambda = 1$ in Theorem 2.2 we have:

Corollary 2.2. $f \in \overline{\mathcal{HM}}(\beta, 1, k, \nu) = \left\{ f : \operatorname{Re} \left((1-k)\frac{(\Omega^\nu f(z))'}{z'} + k\frac{(\Omega^\nu f(z))''}{z''} \right) > \beta \right\}$ if and only if

$$\sum_{n=2}^{\infty} n^2(n-1)(1-k+nk)B(n-1, \nu)|a_n| + \sum_{n=1}^{\infty} n^2(n-1)|nk+k-1|B(n-1, 2-\nu)|b_n| < 1-\beta.$$

Putting $k = 1$ in Theorem 2.2 we have:

Corollary 2.3. $f \in \overline{\mathcal{HM}}(\beta, \lambda, 1, \nu) = \left\{ f : \operatorname{Re} \left((1-\lambda)\frac{\Omega^\nu f(z)}{z} + \lambda\frac{(\Omega^\nu f(z))'}{z'} \right) > \beta \right\}$ if and only if

$$\sum_{n=2}^{\infty} n(n-1)[1+\lambda(n^2-1)]B(n-1, 2-\nu)(|a_n| + |b_n|) < 1-\beta.$$

Finally putting $k = 0$ in Theorem 2.2 we obtain:

Corollary 2.4. $f \in \overline{\mathcal{HM}}(\beta, \lambda, 0, \nu) = \left\{ f : \operatorname{Re} \left((1-\lambda)\frac{\Omega^\nu f(z)}{z} + \lambda\frac{(\Omega^\nu f(z))'}{z'} \right) > \beta \right\}$ if and only if

$$\sum_{n=2}^{\infty} n(n-1)[1+\lambda(n-1)]B(n-1, 2-\nu)|a_n| + \sum_{n=1}^{\infty} n(n-1)|1-\lambda(n+1)|B(n-1, 2-\nu)|b_n| < 1-\beta.$$

Theorem 2.3. $f \in \overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ if and only if

$$(2.5) \quad f(z) = t_1 z + \sum_{n=2}^{\infty} t_n f_n(z) + \sum_{n=1}^{\infty} s_n g_n(z) \quad (z \in \mathcal{D}),$$

where $t_i \geq 0$, $s_i \geq 0$, $t_1 + \sum_{n=2}^{\infty} t_n + \sum_{n=1}^{\infty} s_n = 1$ and

$$f_n(z) = z - \frac{1-\beta}{\phi(n, k, \lambda, \nu)} z^n,$$

$$g_n(z) = z + \frac{1-\beta}{|\psi(n, k, \lambda, \nu)|} \bar{z}^n.$$

Proof. Let f be of the form (2.5) then we have

$$\begin{aligned} f(z) &= t_1 z + \sum_{n=2}^{\infty} t_n \left(z - \frac{1-\beta}{\phi(n, k, \lambda, \nu)} z^n \right) + \sum_{n=1}^{\infty} s_n \left(z + \frac{1-\beta}{|\psi(n, k, \lambda, \nu)|} \bar{z}^n \right) \\ &= z - \sum_{n=2}^{\infty} \frac{1-\beta}{\phi(n, k, \lambda, \nu)} t_n z^n + \sum_{n=1}^{\infty} \frac{1-\beta}{|\psi(n, k, \lambda, \nu)|} s_n \bar{z}^n. \end{aligned}$$

Therefore we have

$$\begin{aligned} &\sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) \frac{1-\beta}{\phi(n, k, \lambda, \nu)} t_n + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| \frac{1-\beta}{|\psi(n, k, \lambda, \nu)|} s_n \\ &= (1-\beta) \left[\sum_{n=2}^{\infty} t_n + \sum_{n=1}^{\infty} s_n \right] = (1-\beta)(1-t_1) \\ &< 1-\beta. \end{aligned}$$

This shows that $f \in \overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$. Conversely suppose that $f \in \overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ letting

$$t_1 = 1 - \sum_{n=2}^{\infty} t_n - \sum_{n=1}^{\infty} s_n,$$

where

$$t_n = \frac{\phi(n, k, \lambda, \nu)}{1 - \beta} |a_n|, \quad s_n = \frac{|\psi(n, k, \lambda, \nu)|}{1 - \beta} |b_n|.$$

We obtain

$$\begin{aligned} f(z) &= z - \sum_{n=2}^{\infty} |a_n| z^n + \sum_{n=1}^{\infty} |b_n| \bar{z}^n \\ &= z - \sum_{n=2}^{\infty} \frac{1 - \beta}{\phi(n, k, \lambda, \nu)} t_n z^n + \sum_{n=1}^{\infty} \frac{1 - \beta}{|\psi(n, k, \lambda, \nu)|} s_n \bar{z}^n. \\ &= z - \sum_{n=2}^{\infty} (z - f_n(z)) t_n + \sum_{n=1}^{\infty} (g_n(z) - z) s_n \\ &= \left(1 - \sum_{n=2}^{\infty} t_n - \sum_{n=1}^{\infty} s_n \right) z + \sum_{n=2}^{\infty} t_n f_n(z) + \sum_{n=1}^{\infty} s_n g_n(z) \\ &= t_1 z + \sum_{n=2}^{\infty} t_n f_n(z) + \sum_{n=1}^{\infty} s_n g_n(z). \end{aligned}$$

This completes the proof. $\square$

3. CONVOLUTION AND CONVEX COMBINATIONS

In the present section we investigate the convolution properties of the class $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$. The convolution of two harmonic function f_1 and f_2 given by

$$(3.1) \quad f_1(z) = z - \sum_{n=2}^{\infty} |a_n| z^n + \sum_{n=1}^{\infty} |b_n| \bar{z}^n, \quad f_2(z) = z - \sum_{n=2}^{\infty} |c_n| z^n + \sum_{n=1}^{\infty} |d_n| \bar{z}^n,$$

is defined by

$$(3.2) \quad (f_1 * f_2)(z) = z - \sum_{n=2}^{\infty} |a_n c_n| z^n + \sum_{n=1}^{\infty} |b_n d_n| \bar{z}^n.$$

Theorem 3.1. *For $0 \leq \beta < \alpha < 1$ let f_1, f_2 be of the form (3.1) such that for every n , $|c_n| < 1$, $|d_n| < 1$. If $f_1, f_2 \in \overline{\mathcal{HM}}(\alpha, \lambda, k, \nu)$ then*

$$f_1 * f_2 \in \overline{\mathcal{HM}}(\alpha, \lambda, k, \nu) \subset \mathcal{HM}(\beta, \lambda, k, \nu).$$

Proof. Considering (3.2) we have

$$\begin{aligned} &\sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n c_n| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n d_n| \\ &< \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_n| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_n| \\ (3.3) \quad &< 1 - \alpha, \end{aligned}$$

and the proof is complete. $\square$

In the last theorem we examine the convex combination properties of the elements of $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$.

Theorem 3.2. *The class $\overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ is closed under convex combination.*

Proof. Suppose that

$$f_i(z) = z - \sum_{n=2}^{\infty} |a_{n,i}| z^n + \sum_{n=1}^{\infty} |b_{n,i}| \bar{z}^n, \quad i = 1, 2, \dots$$

then the convex combinations of f_i may be written as

$$\sum_{i=1}^{\infty} t_i f_i(z) = z - \sum_{n=2}^{\infty} \left(\sum_{i=1}^{\infty} t_i |a_{n,i}| \right) z^n + \sum_{n=1}^{\infty} \left(\sum_{i=1}^{\infty} t_i |b_{n,i}| \right) \bar{z}^n,$$

where $\sum_{i=1}^{\infty} t_i = 1$, $0 \leq t_i \leq 1$. Since

$$\sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_{n,i}| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_{n,i}| < 1 - \beta,$$

so we have

$$\begin{aligned} & \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) \left(\sum_{i=1}^{\infty} t_i |a_{n,i}| \right) + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| \left(\sum_{i=1}^{\infty} t_i |b_{n,i}| \right) \\ &= \sum_{i=1}^{\infty} t_i \left\{ \sum_{n=2}^{\infty} \phi(n, k, \lambda, \nu) |a_{n,i}| + \sum_{n=1}^{\infty} |\psi(n, k, \lambda, \nu)| |b_{n,i}| \right\} \\ &< (1 - \beta) \sum_{i=1}^{\infty} t_i = 1 - \beta. \end{aligned}$$

This shows that $\sum_{i=1}^{\infty} t_i f_i(z) \in \overline{\mathcal{HM}}(\beta, \lambda, k, \nu)$ and the proof is complete. $\square$

REFERENCES

- [1] O. P. Ahuja and J. M. Jahangiri, On a linear combination of classes of multivalently harmonic functions, *Kyungpook Math. J.* **12** (2002), 61-70.
- [2] S. s. Bhoosnurmath and S. R. Swamay, Certain classes of analytic functions with negative coefficients, *Indian J. Math.* **27** (1986), 89-100.
- [3] J. Clunie and T. Sheil Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. A. I. Math.* **9** (1984), 25-40.
- [4] A. Ebadian and A. Tehranchi, On certain class of harmonic p -valent functions by applying the Ruscheweyh derivatives, to appear in *Filomat*.
- [5] J. M. Jahangiri, Harmonic functions starlike in the unit disk, *J. Math. Anal. Appl.* **235** (1999), 470-477.
- [6] A. Lewy, On the non-vanishing of the Jacobian in certain one-to-one mapping, *Bull. Amer. Math. soc.*, **42** (1936), 689-710.
- [7] S. Owa, On the distortion theorems, I, *Kyungpook Math. J.* **18** (1978), 53-59.
- [8] S. Owa and H. M. Srivastava, Univalent and starlike generalized hypergeometric functions, *Canada. J. Math.* **39** (1987), 1057-1072.
- [9] H. M. Srivastava and S. Owa, An application of fractional derivative, *Math. Japon.* **29** (1981), 383-389.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SEMNAN UNIVERSITY, SEMNAN, IRAN
E-mail address: madjid.eshaghi@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN
E-mail address: sa40shams@yahoo.com
E-mail address: a.ebadian@mail.urmia.ac.ir